NEW YORK SMILE INSTITUTE AESTHETIC & IMPLANT DENTISTRY



I. INSTRUCTIONS:

- Make sure the "bite" is correct and you are not hitting too high on the treated area. All teeth should hit at ths same time. If you feel there needs to be an adjustment please make an appointment immediately!
- AVOID eating anything to hard or very cold on the treated area for about 5 days unless otherwise instructed.
- Experiencing slight Hot and Cold sensitivity is normal and should be expected.
- In case of bleeding, place gauze over bleeding area and close mouth to bring pressure to affected area. Let the gauze remain in place with the mouth closed for at least 20 minutes. In addition change cotton gauze EVERY 20 minutes.
- If you are going on any long vacations or trips longer than 3 weeks, please let the doctor know as soon as possible.
- Rinse your mouth thoroughly after meals and before going to bed with **Hydrogen Peroxide & Natural Dentist Rinse** mixed with a glass of warm water.
- Brush teeth as usual. You **CANNOT** Floss, the Temporary Veneers are connected like a Bridge.
- Use a proxybrush between the teeth if it has been prescribed

II. AVOID THESE FOODS FOR THE FIRST 2 WEEKS:

Bagels	Popcorn	Taffy
Ice	Nuts	Sticky Candies
Pretzels	Crusty Food	Chewing Gum

III. FOR MORE THAN A 4 TOOTH TEMPORARY BRIDGE: Best Foods for Nutrition

Pastas	Puddings	Mashed Foods	Boiled Fish	Frozen Yogurt
Rice	Bananas	Steamed Foods	Apple Sause	Milk Shakes